# Waters™

Application Note

## Penicillin G in Pork

Waters Corporation



This is an Application Brief and does not contain a detailed Experimental section.

## Abstract

This method can be used to monitor Penicillin G in pork.

### Introduction

World organizations are concerned about the overuse of antibiotics and antibacterials levels in foods due to possible bacteria resistances and health concerns.


## Experimental

#### Pre-treatment

- 1. Homogenize 3 g of sample with 3 mL of 5% sodium tungstate, 3 mL of 0.17 M sulfuric acid and 30 mL of water.
- 2. Centrifuge 3100 rpm for 10 minutes.
- 3. Filter with glass fiber filter.

#### **SPE Procedure**

## Sep-Pak® Plus Short $C_{18}$ , 360 mg



#### LC Conditions

System: ACQUITY UPLC

Column: ACQUITY UPLC BEH  $C_{18}$ , 1.7  $\mu$ m, 2.1  $\times$  50 mm

Flow rate: 600  $\mu$ L/min

Mobile phase A: 0.1% formic acid in water

Mobile phase B: 0.1% formic acid in acetonitrile

#### Gradient

| Time<br>(min) | %A | %B |
|---------------|----|----|
| 0             | 90 | 10 |
| 5             | 10 | 90 |
| 5.5           | 90 | 10 |
| 6             | 90 | 10 |

#### **MS Conditions**

MS System: Waters Quattro Premier XE


Ionization mode: Positive electrospray (ESI+)

Multiple reaction monitoring

MRM transitions: 1: 335160

2:335176

## Results and Discussion



Representative chromatogram for 5 ppb penicillin spiked in pork muscle.

| Compound Name: penicillin MRM 335→176 | RT    | Area  |
|---------------------------------------|-------|-------|
| pork blank1                           | 1.83  | 4.50  |
| pork blank2                           | 1.83  | 3.23  |
| 5 ppb spiked pork 1                   | 1.83  | 23.30 |
| 5 ppb spiked pork 2                   | 1.82  | 18.54 |
| 5 ppb spiked pork 3                   | 1.83  | 21.82 |
| 5 ppb spiked pork 4                   | 1.83  | 15.45 |
| 5 ppb spiked pork 5                   | 1.82  | 21.50 |
| RSD (%)                               | 15.56 |       |
| Recovery (%)                          | 58.26 |       |

| Compound Name: penicillin MRM 335→160 | RT    | Area  |
|---------------------------------------|-------|-------|
| pork blank1                           | 1.82  | 5.81  |
| pork blank2                           | 1.83  | 5.02  |
| 5 ppb spiked pork 1                   | 1.83  | 42.97 |
| 5 ppb spiked pork 2                   | 1.83  | 26.59 |
| 5 ppb spiked pork 3                   | 1.83  | 21.89 |
| 5 ppb spiked pork 4                   | 1.82  | 25.22 |
| 5 ppb spiked pork 5                   | 1.83  | 39.03 |
| RSD (%)                               | 29.75 |       |
| Recovery (%)                          | 78.30 |       |

Recovery results for 5 ppb penicillin spiked in pork muscle.

## Featured Products

ACQUITY UPLC System <a href="https://www.waters.com/514207">https://www.waters.com/514207</a>

720002592, April 2008

© 2021 Waters Corporation. All Rights Reserved.