Waters™



ACQUITY UPLC Analysis of Water Soluble Vitamins

Waters Corporation



This is an Application Brief and does not contain a detailed Experimental section.

Abstract

This application brief demonstrates the UPLC Analysis of water soluble vitamins.

Introduction

Structures

Nicotinamide

Pyridoxal

Nicotinic acid

d Ascorbic acid

Riboflavin

Thiamine

B12

Compounds

- 1. Nicotinamide (25 μ g/mL)
- 2. Pyridoxal (50 μg/mL)
- 3. Riboflavin (50 μ g/mL)
- 4. Nicotinic acid (50 μg/mL)

5. Inlamine (50 μg/mL)	
6. Ascorbic acid (25 μg/mL)	
7. B12 (50 μg/mL)	
8. Folic Acid (25 μg/mL)	
Experimental	
Test conditions	
Column:	ACQUITY UPLC BEH Amide, 2.1 x 50 mm, 1.7 μm
Part Number:	186004800
Mobile Phase A:	50/50 MeCN/H $_2$ O with 10 mM CH $_3$ COONH $_4$ and 0.04 % NH $_4$ OH, pH 9.0
Mobile Phase B:	90/10 MeCN/H $_2$ O with 10 mM CH $_3$ COONH $_4$ and 0.04 % NH $_4$ OH, pH 9.0
Flow Rate:	0.5 mL/min
Injection Volume:	5 μL (PLNO)
Sample Diluent:	75/25 MeCN/MeOH with 0.2% HCOOH
Column Temperature:	30 °C
Weak Needle Wash:	95/5 MeCN/H ₂ O
Detection:	UV @ 265nm
Sampling Rate:	20 points/sec

Filter Time Constant:

0.2


Instrument:

Waters ACQUITY UPLC with ACQUITY UPLC PDA Detector

Gradient:

Time (min)	%A	%B
Initial	0.1	99.9
3.50	70.0	30.0
3.51	0.1	99.9
7.50	0.1	99.9

Results and Discussion



Featured Products

- ACQUITY UPLC System https://www.waters.com/514207
- ACQUITY UPLC PDA Detector https://www.waters.com/514225

WA60093, June 2009

©2019 Waters Corporation. All Rights Reserved.